arXiv:1203.3290v1 [math.DG] 15 Mar 2012

ALMOST CONTACT B-METRIC MANIFOLDS WITH CURVATURE TENSORS OF KÄHLER TYPE

MANCHO MANEV, MIROSLAVA IVANOVA

Abstract. On 5-dimensional almost contact B-metric manifolds, the form of any φ-Kähler-type tensor (i.e. a tensor satisfying the properties of the curvature tensor of the Levi-Civita connection in the special class of the parallel structures on the manifold) is determined. The associated 1-forms are derived by the scalar curvatures of the φ-Kähler-type tensor for the φ-canonical connection on the manifolds from the main classes with closed 1-forms.

Introduction

The curvature properties of the almost contact B-metric manifolds are investigated with respect to the Levi-Civita connection ∇ and other linear connection preserving the structures of the manifold. A significant role play such connections, which curvature tensors possess the properties of the curvature tensor of ∇ in the class with ∇-parallel structures.

The present paper[1] is organized as follows. In Sec. 1, we give some necessary facts about the considered manifolds. Sec. 2 is devoted to the φ-Kähler-type tensors, i.e. the tensors satisfying the properties of the curvature tensor of ∇ in the special class $\mathcal{F}_0$. In Sec. 3, it is determined the form of any φ-Kähler-type tensor L on a 5-dimensional manifold under consideration. In Sec. 4, it is proved that the associated 1-forms θ

2000 *Mathematics Subject Classification.* 53C05, 53C15, 53C50.

Key words and phrases. Almost contact manifold; B-metric; natural connection; canonical connection; Kähler-type tensor; totally real 2-plane; sectional curvature; scalar curvature.

[1] Partially supported by project NI11-FMI-004 of the Scientific Research Fund, Paisii Hilendarski University of Plovdiv, Bulgaria

and θ^* are derived by the non-φ-holomorphic pair of scalar curvatures of the φ-Kähler-type tensor for the φ-canonical connection on the manifolds from the main classes with closed 1-forms. In Sec. 5, some of the obtained results are illustrated by a known example.

1. PRELIMINARIES

Let $(M, \varphi, \xi, \eta, g)$ be an almost contact manifold with B-metric or an *almost contact B-metric manifold*, i.e. M is a $(2n + 1)$-dimensional differentiable manifold with an almost contact structure (φ, ξ, η) consisting of an endomorphism φ of the tangent bundle, a vector field ξ, its dual 1-form η as well as M is equipped with a pseudo-Riemannian metric g of signature $(n, n + 1)$, such that the following relations are satisfied

$$\varphi\xi = 0, \quad \varphi^2 = -\operatorname{Id} + \eta \otimes \xi, \quad \eta \circ \varphi = 0, \quad \eta(\xi) = 1,$$
$$g(x, y) = -g(\varphi x, \varphi y) + \eta(x)\eta(y)$$

for arbitrary x, y of the algebra $\mathfrak{X}(M)$ on the smooth vector fields on M.

Further, x, y, z will stand for arbitrary elements of $\mathfrak{X}(M)$.

The associated metric $\tilde{g}$ of g on M is defined by

$$\tilde{g}(x, y) = g(x, \varphi y) + \eta(x)\eta(y).$$

Both metrics g and $\tilde{g}$ are necessarily of signature $(n, n+1)$. The manifold $(M, \varphi, \xi, \eta, \tilde{g})$ is also an almost contact B-metric manifold.

The structural tensor F of type (0,3) on $(M, \varphi, \xi, \eta, g)$ is defined by $F(x, y, z) = g\big((\nabla_x \varphi)\, y, z\big)$. It has the following properties:

$$F(x, y, z) = F(x, z, y) = F(x, \varphi y, \varphi z) + \eta(y)F(x, \xi, z) + \eta(z)F(x, y, \xi).$$

The following 1-forms are associated with F:

$$\theta(z) = g^{ij}F(e_i, e_j, z), \quad \theta^*(z) = g^{ij}F(e_i, \varphi e_j, z), \quad \omega(z) = F(\xi, \xi, z),$$

where g^{ij} are the components of the inverse matrix of g with respect to a basis $\{e_i; \xi\}$ $(i = 1, 2, \dots, 2n)$ of the tangent space T_pM of M at an arbitrary point $p \in M$. Obviously, the equality $\omega(\xi) = 0$ and the relation $\theta^* \circ \varphi = -\theta \circ \varphi^2$ are always valid.

A classification of the almost contact manifolds with B-metric with respect to F is given in [3]. This classification includes eleven basic classes $\mathcal{F}_1$, $\mathcal{F}_2$, …, $\mathcal{F}_{11}$. Their intersection is the special class $\mathcal{F}_0$ determined by

$F(x,y,z)=0$. Hence $\mathcal{F}_0$ is the class of almost contact B-metric manifolds with ∇-parallel structures, i.e. $\nabla\varphi=\nabla\xi=\nabla\eta=\nabla g=\nabla\tilde{g}=0$.

In the present paper we consider the manifolds from the so-called main classes $\mathcal{F}_1$, $\mathcal{F}_4$, $\mathcal{F}_5$ and $\mathcal{F}_{11}$, shortly the $\mathcal{F}_i$-manifolds ($i=1,4,5,11$). These classes are the only classes where the tensor F is expressed by the metrics g and $\tilde{g}$. They are defined as follows:

$$\begin{aligned}
\mathcal{F}_1: &\quad F(x,y,z)=\frac{1}{2n}\big\{g(x,\varphi y)\theta(\varphi z)+g(\varphi x,\varphi y)\theta(\varphi^2 z)\big\}_{(y\leftrightarrow z)};\\
\mathcal{F}_4: &\quad F(x,y,z)=-\frac{\theta(\xi)}{2n}\big\{g(\varphi x,\varphi y)\eta(z)+g(\varphi x,\varphi z)\eta(y)\big\};\\
\mathcal{F}_5: &\quad F(x,y,z)=-\frac{\theta^*(\xi)}{2n}\big\{g(x,\varphi y)\eta(z)+g(x,\varphi z)\eta(y)\big\};\\
\mathcal{F}_{11}: &\quad F(x,y,z)=\eta(x)\left\{\eta(y)\omega(z)+\eta(z)\omega(y)\right\},
\end{aligned}$$

where (for the sake of brevity) we use the denotation $\{A(x,y,z)\}_{(y\leftrightarrow z)}$ — instead of $\{A(x,y,z)+A(x,z,y)\}$ for any tensor $A(x,y,z)$.

Let us remark that the class $\mathcal{F}_1\oplus\mathcal{F}_4\oplus\mathcal{F}_5\oplus\mathcal{F}_{11}$ is the odd-dimensional analogue of the class $\mathcal{W}_1$ of the conformal Kähler manifolds of the almost complex manifold with Norden metric, introduced in [4].

2. Curvature-like tensors

Let $R=[\nabla,\nabla]-\nabla_{[\ ,\ ]}$ be the curvature (1,3)-tensor of the Levi-Civita connection ∇. We denote the curvature $(0,4)$-tensor by the same letter: $R(x,y,z,w)=g(R(x,y)z,w)$.

The Ricci tensor ρ and the scalar curvature τ for R as well as their associated quantities are defined respectively by

$$\begin{aligned}
\rho(y,z)&=g^{ij}R(e_i,y,z,e_j), &\qquad \tau&=g^{ij}\rho(e_i,e_j),\\
\rho^*(y,z)&=g^{ij}R(e_i,y,z,\varphi e_j), &\qquad \tau^*&=g^{ij}\rho^*(e_i,e_j).
\end{aligned}$$

Definition 2.1 ([12]). Each (0,4)-tensor L on (M,φ,ξ,η,g) having the following properties is called a *curvature-like tensor*:

$$L(x,y,z,w)=-L(y,x,z,w)=-L(x,y,w,z), \tag{2.1}$$

$$\mathop{\mathfrak{S}}_{x,y,z} L(x,y,z,w)=0. \tag{2.2}$$

The above properties are a characteristic of the curvature tensor R.

Similarly of (2), the Ricci tensor, the scalar curvature and their associated quantities are determined for each curvature-like tensor L.

Definition 2.2 ([12]). A curvature-like tensor L on (M,φ,ξ,η,g) is called a *φ-Kähler-type tensor* if it satisfies the condition

$$L(x,y,\varphi z,\varphi w)=-L(x,y,z,w). \tag{2.3}$$

This property is a characteristic of R on a $\mathcal{F}_0$-manifold. Moreover, (2.3) is similar to the property for a Kähler-type tensor with respect to J on an almost complex manifold with Norden metric ([1]).

Lemma 2.1. *If L is a φ-Kähler-type tensor on (M,φ,ξ,η,g), then the following properties are valid:*

$$L(\varphi x,\varphi y,z,w)=L(x,\varphi y,\varphi z,w)=-L(x,y,z,w), \tag{2.4}$$

$$L(\xi,y,z,w)=L(x,\xi,z,w)=L(x,y,\xi,w)=L(x,y,z,\xi)=0, \tag{2.5}$$

$$L(\varphi x,y,z,w)=L(x,\varphi y,z,w)=L(x,y,\varphi z,w)=L(x,y,z,\varphi w). \tag{2.6}$$

Proof. The equalities (2.4) and (2.5) follow immediately from (2.1), (2.2) and (2.3). The properties (2.4) and (2.5) imply (2.6). □

We consider an associated tensor L^* of L by the equality

$$L^*(x,y,z,w)=L(x,y,z,\varphi w).$$

Let us remark, the tensor L^* is not a curvature-like tensor at all. If L is a φ-Kähler-type tensor, then L^* is also a φ-Kähler-type tensor. Then the properties in Lemma 2.1 are valid for L^*. Obviously, the associated tensor of L^*, i.e. $(L^*)^*$, is $-L$. Consequently, we have the following

Corollary 2.2. *Let L and its associated tensor L^* be φ-Kähler-type tensors on (M,φ,ξ,η,g). Then we have*

$$\begin{aligned} \rho(L^*)&=\rho^*(L), \qquad & \rho^*(L^*)&=-\rho(L),\\ \tau(L^*)&=\tau^*(L), \qquad & \tau^*(L^*)&=-\tau(L). \end{aligned}$$

2.1. **Examples of curvature-like tensors of φ-Kähler type.** Let us consider the following basic tensors of type (0,4) derived by the structural tensors of (M,φ,ξ,η,g) and an arbitrary tensor S of type (0,2):

$$\begin{aligned}
\psi_1(S)(x,y,z,w) &= \big\{g(y,z)S(x,w)+g(x,w)S(y,z)\big\}_{[x\leftrightarrow y]},\\
\psi_2(S)(x,y,z,w) &= \big\{g(y,\varphi z)S(x,\varphi w)+g(x,\varphi w)S(y,\varphi z)\big\}_{[x\leftrightarrow y]},\\
\psi_3(S)(x,y,z,w) &= -\big\{g(y,z)S(x,\varphi w)+g(y,\varphi z)S(x,w)\\
&\qquad +g(x,\varphi w)S(y,z)+g(x,w)S(y,\varphi z)\big\}_{[x\leftrightarrow y]},\\
\psi_4(S)(x,y,z,w) &= \big\{\eta(y)\eta(z)S(x,w)+\eta(x)\eta(w)S(y,z)\big\}_{[x\leftrightarrow y]},\\
\psi_5(S)(x,y,z,w) &= \big\{\eta(y)\eta(z)S(x,\varphi w)+\eta(x)\eta(w)S(y,\varphi z)\big\}_{[x\leftrightarrow y]},
\end{aligned}$$

where we use the denotation $\{A(x,y,z)\}_{[x\leftrightarrow y]}$ instead of the difference $A(x,y,z)-A(y,x,z)$ for any tensor $A(x,y,z)$. The tensor $\psi_1(S)$ coincides with the known Kulkarni-Nomizu product of the tensors g and S.

The five tensors $\psi_i(S)$ are not curvature-like tensors at all. In [12] and [9], it is proved that on an almost contact B-metric manifold:

1. $\psi_1(S)$ and $\psi_4(S)$ are curvature-like tensors if and only if $S(x,y)=S(y,x)$;
2. $\psi_2(S)$ and $\psi_5(S)$ are curvature-like tensors if and only if $S(x,\varphi y)=S(y,\varphi x)$;
3. $\psi_3(S)$ is a curvature-like tensor if and only if $S(x,y)=S(y,x)$ and $S(x,\varphi y)=S(y,\varphi x)$.

Moreover, both of the tensors $\psi_1(S)-\psi_2(S)-\psi_4(S)$ and $\psi_3(S)+\psi_5(S)$ are of φ-Kähler type if and only if the tensor S is symmetric and hybrid with respect φ, i.e. $S(x,y)=S(y,x)$ and $S(x,y)=-S(\varphi x,\varphi y)$. In this case, their associated tensors are the following:

$$\begin{aligned}
&(\psi_1-\psi_2-\psi_4)^*(S)=-(\psi_3+\psi_5)(S),\\
&(\psi_3+\psi_5)^*(S)=(\psi_1-\psi_2-\psi_4)(S).
\end{aligned}$$

The following tensors π_i $(i=1,2,\dots,5)$, derived only by the metric tensors of (M,φ,ξ,η,g), play an important role in differential geometry of an almost contact B-metric manifold:

$$\pi_i = \frac{1}{2}\psi_i(g),\ (i=1,2,3); \qquad \pi_i = \psi_i(g),\ (i=4,5).$$

In [12], it is proved that π_i $(i=1,2,\dots,5)$ are curvature-like tensors and the tensors

$$L_1 = \pi_1 - \pi_2 - \pi_4, \qquad L_2 = \pi_3 + \pi_5$$

are φ-Kähler-type tensors. Their associated φ-Kähler-type tensors are as follows

$$L_1^* = -L_2, \qquad L_2^* = L_1.$$

3. φ-KÄHLER-TYPE TENSORS ON A 5-DIMENSIONAL ALMOST CONTACT B-METRIC MANIFOLD

Let α be a non-degenerate totally real section in T_pM, $p \in M$, and α be orthogonal to ξ with respect to g, i.e. $\alpha \perp \varphi\alpha$, $\alpha \perp \xi$. Let $k(\alpha;p)(L)$ and $k^*(\alpha;p)(L)$ be the scalar curvatures of α with respect to a curvature-like tensor L, i.e.

$$k(\alpha;p)(L) = \frac{L(x,y,y,x)}{\pi_1(x,y,y,x)}, \qquad k^*(\alpha;p)(L) = \frac{L(x,y,y,\varphi x)}{\pi_1(x,y,y,x)},$$

where $\{x,y\}$ is an arbitrary basis of α.

We recall two known propositions for constant sectional curvatures.

Theorem 3.1 ([16]). *Let (M,φ,ξ,η,g) $(\dim M \geq 5)$ be an almost contact B-metric $\mathcal{F}_0$-manifold. Then M is of constant totally real sectional curvatures $\nu = \nu(p)(R) = k(\alpha;p)(R)$ and $\nu^* = \nu^*(p)(R) = k^*(\alpha;p)(R)$ if and only if $R = \nu L_1 + \nu^* L_2$. Both functions ν and ν^* are constant if M is connected and* $\dim M \geq 7$.

Theorem 3.2 ([17]). *Each 5-dimensional almost contact B-metric $\mathcal{F}_0$-manifold has point-wise constant sectional curvatures*

$$\nu(p)(R) = k(\alpha;p)(R), \qquad \nu^*(p)(R) = k^*(\alpha;p)(R).$$

In this relation, we give the following

Theorem 3.3. *Let (M,φ,ξ,η,g) be a 5-dimensional almost contact B-metric manifold. Then each φ-Kähler-type tensor has the form*

$$L = \nu L_1 + \nu^* L_2,$$

where $\nu=\nu(L)$ and $\nu^*=\nu^*(L)=\nu(L^*)$ are the sectional curvatures of the totally real 2-planes orthogonal to ξ in T_pM, $p\in M$, with respect to L. Moreover, (M,φ,ξ,η,g) is of point-wise contact sectional curvatures of the totally real 2-planes orthogonal to ξ with respect to L.

Proof. Let $\{e_1,e_2,\varphi e_1,\varphi e_2,\xi\}$ be an adapted φ-basis of T_pM with respect to g, i.e.

$$-g(e_1,e_1)=-g(e_2,e_2)=g(\varphi e_1,\varphi e_1)=g(\varphi e_2,\varphi e_2)=1,$$
$$g(e_i,\varphi e_j)=0,\quad \eta(e_i)=0\quad (i,j\in\{1,2\}).$$

Then an arbitrary vector in T_pM has the form $x=x^1e_1+x^2e_2+\tilde{x}^1\varphi e_1+\tilde{x}^2\varphi e_2+\eta(x)\xi$. Using properties (2.1), (2.2) and (2.3) for $L(x,y,z,w)$, we obtain immediately $L=\nu L_1+\nu^*L_2$, where $\nu=L(e_1,e_2,e_2,e_1)$, $\nu^*=L(e_1,e_2,e_2,\varphi e_1)=\nu(L^*)=L^*(e_1,e_2,e_2,e_1)$ are the sectional curvatures of α with respect to L, because $\pi_1(e_1,e_2,e_2,e_1)=1$.

Then, if $\{x,y\}$ is an adapted φ-basis of an arbitrary totally real 2-plane α orthogonal to ξ, i.e.

$$g(x,y)=g(x,\varphi x)=g(x,\varphi y)=g(y,\varphi y)=\eta(x)=\eta(y)=0,$$

we get $k(\alpha;p)(L)=\nu(p)(L)$, $k^*(\alpha;p)(L)=\nu^*(p)(L)$, taking into account the expression $L=\nu L_1+\nu^*L_2$. Therefore, (M,φ,ξ,η,g) is of point-wise contact sectional curvatures of α with respect to L. □

The restriction of Theorem 3.3 in $\mathfrak{F}_0$ confirms Theorem 3.1 because R is a φ-Kähler-type tensor on a $\mathfrak{F}_0$-manifold.

3.1. Curvature tensor of a natural connection on a 5-dimensional almost contact B-metric manifold.

In [10], it is introduced the notion of a *natural connection* on the manifold (M,φ,ξ,η,g) as a linear connection D, with respect to which the almost contact structure (φ,ξ,η) and the B-metric g are parallel, i.e. $D\varphi=D\xi=D\eta=Dg=0$. According to [13], a necessary and sufficient condition a linear connection D to be natural on (M,φ,ξ,η,g) is $D\varphi=Dg=0$.

Let K be curvature tensor of a natural connection D with torsion T. Then K satisfies (2.1) and (2.3). Instead of (2.2), we have the following form of the first Bianchi identity ([5])

$$\underset{x,y,z}{\mathfrak{S}} K(x,y,z,w)=\underset{x,y,z}{\mathfrak{S}}\left\{T(T(x,y),z,w)+\left(D_xT\right)(y,z,w)\right\}.$$

If we set the condition $\underset{x,y,z}{\mathfrak{S}} K(x,y,z,w)=0$ as for the curvature tensor R then K is a φ-Kähler-type tensor and satisfies the condition of Theorem 3.3. Therefore, we obtain

Corollary 3.4. *Let (M,φ,ξ,η,g) be a 5-dimensional almost contact B-metric manifold with a natural connection D with curvature tensor K of φ-Kähler-type. Then K has the form*

$$K=\nu L_1+\nu^* L_2,$$

where $\nu=\nu(K)$ and $\nu^*=\nu^*(K)=\nu(K^*)$ are the sectional curvatures of the totally real 2-planes orthogonal to ξ in T_pM, $p\in M$, with respect to K. Moreover, (M,φ,ξ,η,g) is of point-wise contact sectional curvatures of the totally real 2-planes orthogonal to ξ with respect to K.

4. CURVATURE TENSOR OF THE φ-CANONICAL CONNECTION

According to [15], a natural connection D is called a *φ-canonical connection* on the manifold (M,φ,ξ,η,g) if the torsion tensor T of D satisfies the following identity:

$$\begin{aligned}\big\{T(x,y,z)-T(x,\varphi y,\varphi z)-\eta(x)\left\{T(\xi,y,z)-T(\xi,\varphi y,\varphi z)\right\}\\ -\eta(y)\left\{T(x,\xi,z)-T(x,z,\xi)-\eta(x)T(z,\xi,\xi)\right\}\big\}_{[y\leftrightarrow z]}=0.\end{aligned}$$

Let us remark that the restriction the φ-canonical connection D of the manifold (M,φ,ξ,η,g) on the contact distribution $\ker(\eta)$ is the unique canonical connection of the corresponding almost complex manifold with Norden metric, studied in [2].

In [12], it is introduced a natural connection on (M,φ,ξ,η,g), defined by

$$D_xy=\nabla_xy+\frac{1}{2}\big\{(\nabla_x\varphi)\,\varphi y+(\nabla_x\eta)\,y\cdot\xi\big\}-\eta(y)\nabla_x\xi. \tag{4.1}$$

In [14], the connection determined by (4.1) is called a *φB-connection*. It is studied for some classes of (M,φ,ξ,η,g) in [12], [6], [7] and [14]. The φB-connection is the odd-dimensional analogue of the B-connection on the corresponding almost complex manifold with Norden metric, studied for the class $\mathcal{W}_1$ in [1].

In [15], it is proved that the φ-canonical connection and the φB-connection coincide on the almost contact B-metric manifolds in a class containing $\mathcal{F}_1 \oplus \mathcal{F}_4 \oplus \mathcal{F}_5 \oplus \mathcal{F}_{11}$.

According to [12], the necessary and sufficient conditions K to be a φ-Kähler-type tensor in $\mathcal{F}_i$ $(i = 1, 4, 5, 11)$ is the associated 1-forms θ, θ^* and $\omega \circ \varphi$ to be closed. These subclasses we denote by $\mathcal{F}_i^0$ $(i = 1, 4, 5, 11)$.

Bearing in mind the second Bianchi identity

$$\underset{x,y,z}{\mathfrak{S}} \left\{ (D_x K)(y, z) + K(T(x, y), z) \right\} = 0,$$

we compute the scalar curvatures for K determined by

$$\tau(K) = g^{ij} \rho(K)_{ij}, \qquad \tau^*(K) = \tau(K^*) = g^{ij} \varphi_j^k \rho(K)_{ik},$$

where $\rho(K)_{ij}$ is the Ricci tensor of K, and then we get the following

Lemma 4.1. *For* $(M, \varphi, \xi, \eta, g)$ *in* $\mathcal{F}_i^0$ $(i = 1, 4, 5, 11)$*, the relations for the scalar curvatures* $\tau = \tau(K)$ *and* $\tau^* = \tau^*(K)$ *of* K *are:*

$$\mathrm{d}\tau \circ \varphi = -\mathrm{d}\tau^* - \frac{1}{n}\left(\tau\theta + \tau^*\theta^*\right), \quad \mathrm{d}\tau^* \circ \varphi = \mathrm{d}\tau - \frac{1}{n}\left(\tau^*\theta - \tau\theta^*\right). \tag{4.2}$$

Obviously, bearing in mind (4.2), we obtain that the pair (τ, τ^*) on $(M, \varphi, \xi, \eta, g)$ is a φ-holomorphic pair of functions, i.e. $\mathrm{d}\tau = \mathrm{d}\tau^* \circ \varphi$ and $\mathrm{d}\tau^* = -\mathrm{d}\tau \circ \varphi$, if and only if the associated 1-forms θ and θ^* are zero. Such one is the case for the class $\mathcal{F}_{11}$.

The system (4.2) can be solved with respect to θ and θ^* and then

$$\theta = -n\left\{\mathrm{d}f_1 + \mathrm{d}f_2 \circ \varphi\right\}, \qquad \theta^* = n\left\{\mathrm{d}f_1 \circ \varphi - \mathrm{d}f_2\right\}, \tag{4.3}$$

where $f_1 = \arctan(\tau^*/\tau)$, $f_2 = \ln\sqrt{\tau^2 + \tau^{*2}}$.

Let us consider the complex-valued function $h = \tau + i\tau^*$ or in polar form $h = |h|e^{i\alpha}$. Then we have $|h| = \sqrt{\tau^2 + \tau^{*2}}$, $\alpha = \arctan(\tau^*/\tau)$.

Bearing in mind that $\operatorname{Log} h = \ln|h| + i\alpha$, then (4.3) take the following form:

$$\theta = -n\left\{\mathrm{d}\alpha + \mathrm{d}(\ln|h|) \circ \varphi\right\}, \quad \theta^* = n\left\{\mathrm{d}\alpha \circ \varphi - \mathrm{d}(\ln|h|)\right\}. \tag{4.4}$$

So, we obtain the following

Theorem 4.2. *For* $(M, \varphi, \xi, \eta, g)$ *in* $\mathcal{F}_i^0$ $(i = 1, 4, 5)$*, the associated 1-forms* θ *and* θ^* *are derived by the non-*φ*-holomorphic pair of the scalar curvatures* (τ, τ^*) *of the* φ*-Kähler-type tensor* K *for the* φ*-canonical connection* D *by* (4.4).

Corollary 4.3.
For $i=1$

$$\theta=n\left\{\mathrm{d}\alpha\circ\varphi^2-\mathrm{d}(\ln|h|)\circ\varphi\right\},\qquad \theta^*=n\left\{\mathrm{d}\alpha\circ\varphi+\mathrm{d}(\ln|h|)\circ\varphi^2\right\};$$

For $i=4$

$$\theta=-n\mathrm{d}\alpha(\xi)\eta,\qquad \theta^*=0;$$

For $i=5$

$$\theta=0,\qquad \theta^*=-n\mathrm{d}(\ln|h|)(\xi)\eta.$$

5. EXAMPLES OF ALMOST CONTACT MANIFOLDS WITH B-METRIC

Let us consider $\mathbb{R}^{2n+2}=\left\{\left(u^1,\dots,u^{n+1};v^1,\dots,v^{n+1}\right)\ \mid u^i,v^i\in\mathbb{R}\right\}$ as a complex Riemannian manifold with the canonical complex structure J and a metric g, defined by $g(x,x)=-\delta_{ij}\lambda^i\lambda^j+\delta_{ij}\mu^i\mu^j$, where $x=\lambda^i\frac{\partial}{\partial u^i}+\mu^i\frac{\partial}{\partial v^i}$. Identifying the point $p=\left(u^1,\dots,u^{n+1};v^1,\dots,v^{n+1}\right)$ in $\mathbb{R}^{2n+2}$ with its positional vector Z, in [3] it is given a hypersurface S defined by

$$g(Z,JZ)=0,\quad g(Z,Z)=\cosh 2t,\quad t>0.$$

The almost contact structure is determined by the conditions:

$$\xi=\frac{1}{\cosh t}Z,\qquad Jx=\varphi x+\eta(x)J\xi,$$

where $x,\varphi x\in T_pS$ and $J\xi\in(T_pS)^\perp$. Then (S,φ,ξ,η,g) is an almost contact B-metric manifold in the class $\mathcal{F}_5$.

Consequently, we characterize (S,φ,ξ,η,g) by means of [8]. We compute the following quantities for the constructed $\mathcal{F}_5$-manifold:

$$\theta=0,\qquad \eta=\sinh t\mathrm{d}t,\qquad \frac{\xi\theta^*(\xi)}{2n}=-\frac{\theta^{*2}(\xi)}{4n^2}=\frac{1}{\cosh^2 t}. \tag{5.1}$$

In [11], it is given that the 1-form θ^* on a $\mathcal{F}_5$-manifold is closed if and only if $x\theta^*(\xi)=\xi\theta^*(\xi)\eta(x)$. By virtue of (5.1), we establish that (S,φ,ξ,η,g) belongs to the subclass $\mathcal{F}_5^0$, since $\mathrm{d}\theta^*=0$.

The condition for the second fundamental form of the hypersurface S, given in [3], the Gauss equation ([5]) and the flatness of $\mathbb{R}^{2n+2}$ imply the following form of the curvature tensor of ∇

$$R=-\frac{1}{\cosh^2 t}\pi_2.$$

Then, taking into account (5.1) and the form of the curvature tensor K of the φ-canonical connection in $\mathcal{F}_5^0$ ([12])

$$K = R + \frac{\xi\theta^*(\xi)}{2n}\pi_4 + \frac{\theta^{*2}(\xi)}{4n^2}\pi_1,$$

we obtain

$$K = \frac{1}{\cosh t}L_1. \tag{5.2}$$

Since L_1 is a φ-Kähler-type tensor, then K is also a φ-Kähler-type tensor. Therefore, we have

$$\nu(K) = K(e_1, e_2, e_2, e_1) = \frac{1}{\cosh^2 t}, \qquad \nu^*(K) = K^*(e_1, e_2, e_2, e_1) = 0,$$

which confirm Theorem 3.3 and Corollary 3.4.

According to (5.2), the scalar curvatures are

$$\tau(K) = \frac{4n(n-1)}{\cosh^2 t}, \qquad \tau^*(K) = 0.$$

Then, taking into account (5.1), the results for $(S, \varphi, \xi, \eta, g)$ confirm also Lemma 4.1, Theorem 4.2 and Corollary 4.3.

References

[1] G. Ganchev, K. Gribachev, V. Mihova. *B-connections and their conformal invariants on conformally Kaehler manifolds with B-metric*, Publ. Inst. Math. (Beograd) (N.S.), vol. 42(56) (1987), 107–121.

[2] G. Ganchev, V. Mihova. *Canonical connection and the canonical conformal group on an almost complex manifold with B-metric*, Ann. Univ. Sofia Fac. Math. Inform., vol. 81 (1987), 195–206.

[3] G. Ganchev, V. Mihova, K. Gribachev. *Almost contact manifolds with B-metric*, Math. Balkanica (N.S.), vol. 7, no. 3-4 (1993), 261–276.

[4] K. I. Gribachev, D. G. Mekerov, G. D. Djelepov. *Generalized B-manifolds*, Compt. rend. Acad. bulg. Sci., vol. 38, no. 3 (1985), 299–302.

[5] S. Kobayshi, K. Nomizu. *Foundations of differential geometry*, Intersc. Publ., New York, 1969.

[6] M. Manev. *Properties of curvature tensors on almost contact manifolds with B-metric*, Proc. of Jubilee Sci. Session of Vassil Levsky Higher Mil. School, Veliko Tarnovo, vol. 27 (1993), 221–227.

[7] M. Manev. *Contactly conformal transformations of general type of almost contact manifolds with B-metric. Applications*, Math. Balkanica (N.S.), vol. 11, no. 3-4 (1997), 347–357.

[8] M. Manev. *Examples of almost contact manifolds with B-metric of some special classes*. Math. Educ. Math., Proc. 26th Spring Conf. of UBM, Plovdiv, 1997, 153–160 (in Bulgarian).

[9] M. Manev. *On the conformal geometry of almost contact manifolds with B-metric*, Ph.D. Thesis, Plovdiv, 1998. (in Bulgarian)

[10] M. Manev. *A connection with totally skew-symmetric torsion on almost contact manifolds with B-metric*, Int. J. Geom. Methods Mod. Phys., vol. 9, no. 5 (2012), arXiv:1001.3800.

[11] M. Manev, K. Gribachev. *Contactly conformal transformations of almost contact manifolds with B-metric*. Serdica Math. J., vol. 19 (1993), 287–299.

[12] M. Manev, K. Gribachev. *Conformally invariant tensors on almost contact manifolds with B-metric*, Serdica Math. J., vol. 20 (1994), 133–147.

[13] M. Manev, M. Ivanova. *A classification of the torsion tensors on almost contact manifolds with B-metric*, arXiv:1105.5715.

[14] M. Manev, M. Ivanova. *A natural connection on some classes of almost contact manifolds with B-metric*, arXiv:1110.3023.

[15] M. Manev, M. Ivanova. *Canonical-type connection on almost contact manifolds with B-metric*, arXiv:1203.0137.

[16] G. Nakova, K. Gribachev. *Submanifolds of some almost contact manifolds with B-metric with codimension two, I*, Math. Balk., vol. 11 (1997), 255–267.

[17] G. Nakova, M. Manev. *Curvature tensors on some five-dimensional almost contact B-metric manifolds*, Math. Educ. Math., Proc. 32nd Spring Conf. of UBM, Sunny Beach, 2003, 192–197.

M. Manev, M. Ivanova
Department of Algebra and Geometry
Faculty of Mathematics and Informatics
University of Plovdiv
236 Bulgaria Blvd
4003 Plovdiv, Bulgaria
`mmanev@uni-plovdiv.bg, mirkaiv@uni-plovdiv.bg`